

CETAPRED

Sterile Ophthalmic Suspension &
Gamma Irradiated Ophthalmic Ointment

سیٹاپریڈ

Composition:

Ophthalmic Suspension:

Each ml contains:

Sulphacetamide sodium 100 mg
Prednisolone acetate 2 mg
Phenylephrine HCL 1.2 mg with
HPMC.

Ophthalmic Ointment:

Each gram contains:

Sulphacetamide sodium 100 mg
Prednisolone acetate 2 mg

Action:

Sulphacetamide sodium at 10% concentration is a potent bacteriostatic agent, effective against a broad range of pathogens, including staphylococci.

The prednisolone content effectively counters the allergic and inflammatory manifestations of blepharitis.

The Phenylephrine component rapidly whitens the engorged vessels in the eye and lid.

Indications:

Nonpurulent blepharitis and blepharo-conjunctivitis (seborrheal, staphylococcal, allergic), non purulent conjunctivitis (allergic and bacterial). A steroid/anti-infective combination is indicated in ocular inflammation when concurrent use of an antimicrobial is judged necessary.

Contraindications:

Acute herpas simplex (dendritic keratitis), Purulent untreated infections, vaccinia vericella and most other viral diseases of the cornea and conjunctiva, ocular tuberculosis and fungal diseases of the eye. Hypersensitivity to any component.

Warnings:

1. In diseases due to micro-organisms infection may be masked enhanced or activated by the steroids.
2. Extended use may cause increase intra-ocular pressure in susceptible individuals. It is advisable that the intra-ocular pressure should be checked frequently.
3. Various ocular disease and long term use of topical corticosteroids have been known to cause corneal and scleral things. Use of topical corticosteroids in the presence of thin corneal or scleral tissue may lead to perforation.
4. Use with caution in patients with known or suspected sensitivity to sulfonamide. If sensitivity or other unwanted reactions occur, discontinue medication.

5. Should be used with caution in the presence of narrow angle glaucoma.
6. Prolonged use may suppress the host immune response in ocular infection.
7. Use of corticosteroid in the treatment of patients with a history of herpes simplex requires great caution.
8. A significant percentage of staphylococcal isolates is resistant to sulfonamides.

Dosage and Administration:

Ophthalmic suspension:

The recommended dosage is 1 to 2 drops three to four times daily, depending upon the severity of the condition.

In general, during early or acute stages of blepharitis, **Cetapred** produces result most rapidly and most effectively with instillation directly into the eye, with the excess spread on the lid when the condition is confined to the lid, **Cetapred** may be applied directly to the site of the lesion.

Ophthalmic Ointment:

A small amount should be applied in the conjunctival sac three times daily or once or twice at night.

Precaution:

For topical ophthalmic use only.

Protect from light and moisture.

Use only on medical advice.

Once opened use within one month.

Do not touch the tip of the tube.

How supplied:

Ophthalmic Suspension:

Cetapred Ophthalmic Suspension is available in 5ml plastic dropper bottle.

Ophthalmic Ointment:

Cetapred Ophthalmic Ointment is supplied as pack of 3.5 grams in tube.

Storage Conditions:

Store in the cool and dry place at 15°C to 30°C.

Keep out of the reach of children.

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات: دو اکوڑنی، گرمی اور نمی سے بچائیں، عضنی اور خشک جگہ پر رکھیں۔

تمام ادویات کو بچوں کی پہنچ سے دور رکھیں۔

دوا کو کھولنے کے بعد ایک ماہ کے اندر استعمال کر لیں۔

انتباہ:

صرف رجسٹرڈ میڈیکل پریکٹیشنر کے نسخے پر فروخت کے لیے۔



Manufactured by:

Vega Pharmaceuticals (Pvt.) Ltd.

30 Km, Multan Road, Lahore - Pakistan.