

EAROCIN

Sterile Ear Drops
Ofloxacin

ایرو سین
اوفلوکساسن

COMPOSITION:

1. Earocin 0.3%

Each ml of Earocin Ear Drops contains, ofloxacin3.0mg.

2. Earocin 0.6%

Each ml of Earocin Ear Drops contains, ofloxacin6.0mg.

(Vega Specifications)

INDICATIONS:

Earocin (Ofloxacin) is indicated for the treatment of Otitis externa, Otitis media with tympanostomy tube, and Chronic suppurative otitis media infections caused by susceptible strains of microorganisms e.g. Pseudomonas aeruginosa, Staphylococcus aureus, Streptococcus pyogenes, Proteus mirabilis, Streptococcus pneumoniae, Haemophilus influenzae, Moraxella catarrhalis, etc.

CONTRA-INDICATIONS:

Earocin is contra-indicated in patients with a history of hypersensitivity to Ofloxacin, to other Quinolones, or to any of its components.

WARNINGS:

For otic use only. Not for Ophthalmic use. Not for Injection. If an allergic reaction is suspected with Ofloxacin, stop the drug. Serious acute hypersensitivity reactions may require immediate emergency treatment.

PRECAUTIONS:

As with other anti-infectives prolonged use may result in over-growth of non susceptible organisms including Fungi. If the infection is not improved after one-week, cultures should be obtained to guide further treatment. If otorrhea persists after a full course of therapy or if two or more episodes of otorrhea occur within six months, further evaluation is recommended to exclude an underlying condition such as cholesteatoma, foreign body or a tumor.

DRUG INTERACTION:

Specific drug interaction studies have not been conducted with Ofloxacin otic solution.

PREGNANCY:

Ofloxacin has not been shown to have any adverse effects on the developing embryo or fetus at the recommended doses.

NURSING MOTHERS:

Because it is not known whether Ofloxacin is excreted in human milk following topical otic administration, a decision should be made in the interest of the patient.

SIDE EFFECTS:

Rashes, itching, discomfort, ear drainage, discharge,

worsening pain.

DOSAGE & ADMINISTRATION:

Earocin 0.3%: 1. Children: (1 to 12 years of age):

Five drops into the affected ear twice daily for ten days or as directed by the doctor.

2. Adults: 12 years of age and older:

Ten drops into the affected ear twice daily for 14 days or as directed by the doctor.

Earocin 0.6%: 1. Children: (1 to 12 years of age):

Two to Three drops into the affected ear twice daily for ten days or as directed by the doctor.

2. Adults: 12 years of age and older:

Five drops into the affected ear twice daily for 14 days or as directed by the doctor.

STORAGE:

Store below 30°C. Protect from heat and sunlight. Keep out of reach of children.

Presentation:

Earocin Ear Drops 0.3% and 0.6% are available in 5ml sterile plastic dropper bottle.

Precaution:

Use with in four weeks of the first opening.

ایرو سین

اوفلوکساسن 0.6%

خوراک:

بچوں کیلئے (1 سے 12 سال کی عمر تک) 2 سے 3 قطرے متاثرہ کان / کانوں میں

دن میں 2 مرتبہ 10 دن تک ڈالیں یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

بالغوں کیلئے (12 سال یا اس سے اوپر) 5 قطرے متاثرہ کان / کانوں میں دن میں

2 مرتبہ 14 دن تک ڈالیں یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ایرو سین

اوفلوکساسن 0.3%

خوراک:

بچوں کیلئے (1 سے 12 سال کی عمر تک) 5 قطرے متاثرہ کان / کانوں میں

دن میں 2 مرتبہ 10 دن تک ڈالیں یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

بالغوں کیلئے (12 سال یا اس سے اوپر) 10 قطرے متاثرہ کان / کانوں میں دن

میں 2 مرتبہ 14 دن تک ڈالیں یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات: دو کو دھوپ اور گرمی سے بچائیں۔

* ڈگری سنٹی گریڈ سے کم درجہ حرارت پر رکھیں۔

تمام ادویات بچوں کی پہنچ سے دور رکھیں۔

پہلی دفعہ بوتل کھولنے کے بعد دو کو چار ہفتوں تک استعمال کیا جاسکتا ہے۔



Manufactured by:

Vega Pharmaceuticals (Pvt.) Ltd.

30 Km, Multan Road, Lahore, Pakistan.