Mydrex

(Tropicamide) Sterile Ophthalmic Solution



COMPOSITION:

Each ml contains: Tropicamide 10mg. USP Specifications Preservative: Thiomersal.

DESCRIPTION:

Tropicamide is pharmacologically related to the parasympatholytic (anticholinergic) group. It is a white crystalline solid melting at 96°C to 96.5°C. It is slightly soluble in water. Tropicamide is a synthetic compound described chemically as Tropic acid-N-ethyl-N (**g**-picolyl) amide.

ACTIONS:

A mydriatic-cycloplegic. Its action is remarkable in that maximum mydriasis and cycloplegia are obtained within 20 to 25 minutes following instillation in the eye. Complete recovery occurs within 5 to 6 hours without the use of a miotic.

INDICATIONS:

For mydriasis and cycloplegia for diagnostic purposes. For cycloplegic effect in refractions, the 1.0% solution is necessary.

CONTRAINDICATIONS: Contraindicated in narrow angle glaucoma.

WARNINGS: FOR OPHTHALMIC USE ONLY - NOT FOR INJECTION.

PRECAUTIONS: As with any mydriatic, caution should be used when intraocular pressure is high

or unknown or when anterior chamber is shallow. In refractions where prolongation of cycloplegia is desirable, one additional drop only is recommended.

ADVERSE REACTIONS:

Transient stinging may occur when drops are installed. **DOSAGE AND ADMINISTRATION:**

For refraction, 1 to 2 drops of 1.0% solution in eye(s) repeated in 5 minutes. If patient is not seen within 20 to 30 minutes, an additional drop may be instilled to prolong mydriatic effect.

HOW SUPPLIED:

(1.0%) in 15 ml plastic bottle.

STORAGE:

Store below 30°C. Protect from heat and sunlight. Keep out of the reach of children. Use within four weeks after first opening.

> خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔ دواکودھوپ اورگرمی سے بچا 'میں۔ ۳۰ ڈگری بینڈی گریڈ ہے کہ درہ جرارت پر کھیں۔ تمام ادویات بچوں کی بینی سے دور کھیں۔ پہلی دفعہ ہوتا کو لیے کے بعد دواکو چار نفتوں تک استعمال کیا جا سکتا ہے۔



Manufactured by: Vega Pharmaceuticals (Pvt.) Ltd. 30 Km, Multan Road, Lahore, Pakistan.