VEGACROM

(Sterile Eye Drops)
(Sodium Cromoglycate - Tetrahydrozoline Hydrochloride)

ویگا کروم
سوڈیم کروموگلائیکیٹ
ٹیٹراہائیڈروزولین ہائیڈروکلورائیڈ

DESCRIPTION:
Vegacrom sterile ophthalmic solution is offered in a sterile container having approx 15 ml sterile ophthalmic solution.

COMPOSITION:
Each ml contains:
Sodium cromoglycate .....40mg.
Tetrahydrozoline Hydrochloride... 0.5 mg.
(Vega Specifications.)
Vehicle: Hydroxypropyl methyl cellulose.
Preservative: Benzalkonium chloride.

Actions:
Sodium cromoglycate appears to act primarily through a local effect by inhibiting the degranulation of sensitized mast cells occuring after exposure to specific antigens, thus inhibiting the release of histamine and SRS-A (Slow reacting substance of anaphylaxis) from the mast cells.

INDICATIONS:
Vegacrom is a highly effective combination to control the symptoms of burning, itching, tearing and photophobia. Vegacrom is particularly suitable in eye allergic conditions like hay-fever conjunctivitis, acute and chronic allergic conjunctivitis, where a prolonged treatment is needed and where irregular use of steriods is required.

CONTRAINDICATIONS:
Vegacrom should not be used in patients with glaucoma, hypertension, cardiac disorders, hyperthyroidism and diabetes. In the treatment of children, the product containing vasoconstrictor should be used under the direct supervision of an ophthalmologist and should not be used in children under five years. Its safety in pregnancy has not been established.

Vegacrom should not be used in patients with known hypersensitivity to any of its components.

ADVERSE REACTIONS:
Transient stinging, burning sensation, irritation, pupillary dialation may occur after local use of product containing, vasoconstrictor. Due to systemic absorption, some patients may complain of drowsiness, headache, hypertension, cardiac irregularities and hyperglycemia. If the condition persists discontinue the medicine and consult an ophthalmologist.

PRECAUTIONS:
Since sodium cromoglycate has a preventive action, therefore the patients showing positive response should continue the treatment. If the treatment is required to be discontinued, the drug must be withdrawn gradually over a period of one week. Symptoms of conjunctivitis may reappear in case of discontinuation.

As the drug contains vasoconstrictor, it should be used with caution in patients taking antidepressive drugs.

Vegacrom must be kept out of the reach of children as accidentally swallowing may cause sedative effects and in some cases other toxic symptoms may appear.

PATIENT INFORMATION:
1. A transient stinging or burning sensation may occur after instillation, which usually disappears with continued use.
2. Do not wear soft contact lenses during treatment. Lenses may be worn a few hours after discontinuation of drug.
3. Administer the drug at regular intervals.

DOSAGE:
Instill 1 to 2 drops into the affected eye(s) 4 times a day or as directed by the physician.

STORAGE AND INSTRUCTIONS:
Store below 30°C.
Use within 4 weeks after first opening the bottle.
Protect from heat and sunlight.
Keep all medicines out of the reach of children.
Use as directed by the physician.

خوراک: ڈاکٹر کی ہدایات کے مطابق استعمال کریں۔
ہدایات: دوا کو دھوپ اور گرمی سے بچائیں۔
۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔
تمام ادویات بچوں کی پہنچ سے دور رکھیں۔
پہلی دفعہ بوتل کھولنے کے بعد دوا کو چار ہفتوں تک استعمال کیا جا سکتا ہے۔
استعمال سے پہلے اچھی طرح ہلا لیں۔

Manufactured by:
Vega Pharmaceuticals (Pvt.) Ltd.
30km, Multan Road, Lahore-Pakistan.