Vegatob Sterile Ear Drops

Tobramycin & Dexamethasone Sodium Phosphate

ویگاٹوب

ٹوبرامائیسن

ڈیکسامیتھاسون سوڈیم فاسفیٹ

Description:
Vegatob is an antibiotic and steroid combination for otic use.

Composition:
Each ml contains:
Tobramycin.....3.0 mg
Dexamethasone Sodium Phosphate eq. to
Dexamethasone.....1.0 mg
(Vega Specs.)
Preservative:
Benzalkonium Chloride.

CLINICAL PHARMACOLOGY:
Dexamethasone is a corticosteroid with a range of anti-inflammatory properties. Corticosteroids reduce edema, fibrin deposition, capillary dilatation, leukocytes migration, capillary-proliferation, deposition of collagen and scar formation, Dexamethasone has been found exceptionally effective in decreasing granulation tissue which usually causes obstruction in tympanostomy tube. The combination of Dexamethasone with Tobramycin facilitates the physical entry of antibiotic drops in the inflammed ear.

TOBRAMYCIN: Tobramycin is active against susceptible strains of the following microorganism: Staphylococci, including S.auerus and S.epidermidis (coagulase-positive and coagulase-coagulase-negative), including penicillin-resistant strains. Streptococci, including some of the Group A beta-hemolytic species, some non-hemolytic species, and some Streptococcus pneumoniae. Pseudomonas aeruginosa, Escherichia coli, Klebsiella pneumonia, Enterobacter aerogenes, Proteus mirabilis, Morganella morganii, most proteus vulgaris strains, Haemophilus influenzea and H,aegyptius,, Moraxella lacunata, and Acinetobacter calcoaceticus (Herellea vaginacola) and some Neisseria species. Bacterial susceptibility studies demonstrate that in some cases microorganisms resistant to Gentamicin remain susceptible to Tobramycin.

INDICATIONS AND USAGE:
VEGATOB is indicated in steroid-responsive inflammatory otic condition for which a corticosteroid is indicated and where a risk of superficial bacterial otic infection exists.

PRECAUTIONS/WARNINGS:
The possibility of fungal infections of the ear canal should be considered after long term steroid use. Use with in four weeks of the first opening.

ADVERSE REACTIONS:
Adverse reactions have occurred with steroidal / anti-infective combination drugs. The most frequent adverse reactions to topical Tobramycin are localized hypersensitivity, including itching and swelling, These reactions occur in less than 4% of patients.

DOSAGE AND ADMINISTRATION:
Children: (1 to 12 years old):
2-3 drops into the affected ear(s) twice daily.
Adults: (12 years and older):
3-4 drops into the affected ear(s) twice daily or as directed by the physician.

STORAGE AND INSTRUCTIONS:
For otic use only.
On prescription only.
Store below 30° C. Protect from heat and sunlight.
Keep out of the reach of children..

PRESENTATION:
Vegatob Sterile Ear Drops available in 5ml Plastic Dropper bottle.

ہدایات:
بچوں کیلئے: ایک سے بارہ سال کی عمر تک دو سے تین قطرے متاثرہ کان میں دن میں
دو دفعہ یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔
بالغوں کے لیے: بارہ سال یا اس سے اوپر تین سے چار قطرے متاثرہ کان میں
دن میں دو دفعہ یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔
۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔
گرمی اور دھوپ سے بچائیں۔
صرف مستند ڈاکٹر کے نسخہ پر استعمال کریں۔
پہلی دفعہ بوتل کھولنے کے بعد دوا کو چار ہفتوں تک استعمال کیا جا سکتا ہے۔

Manufactured by:
Vega Pharmaceuticals (Pvt.) Ltd.
30 Km, Multan Road, Lahore, Pakistan.